

## Vaccinations: “not just for kids”<sup>(1)</sup>

Jean-Pierre Michel

Department of Rehabilitation and Geriatrics, Geneva Medical School and University Hospitals, Thonex-Geneve, Switzerland

Vaccines which have drastically reduced the burden of childhood diseases are not yet accepted as contributing to improving healthy aging (1-3). The non-respect and/or absence of sustainability in vaccine programs do not enable maintenance of life-long protection against such childhood diseases as diphtheria (4), measles (5) and pertussis (6), which may result in increasing incidence of these diseases in adults in the future (7). Infectious diseases in older people also remain a significant cause of morbidity and mortality in the increasing population of adults aged 60 years and older (8, 9). The high burden of infectious diseases in this segment of the population is disproportionate, as many of these diseases are vaccine-preventable (10).

In 2001, the lower respiratory infections which represented the fourth most frequent cause of death in high-income countries – 4.4% of total deaths (11) – are three times more lethal in old than in young adults (12). Surprisingly, tetanus is still a current active disease, with 210 cases in Portugal between 1993 and 2002 (13), 175 in England between 1984 and 2000 (14) and 158 in Poland between 1998 and 2005 (15, 16) affecting essentially adults over 55 years of age. In the 1990s, a diphtheria epidemic in the newly independent states of the former Soviet Union claimed over 3000 lives, essentially among persons aged 35 to 50 years (4). The notification incidence of pertussis in the Netherlands also increased notably between 1998-2001 and 2002-2005 in the older population group (6). The annual incidence of Herpes zoster in the general population is estimated at 3.6-14.2 cases/1000, and the recurrence risk is multiplied by 8 to 10 in adults over the age of 60 (17, 18).

Without a specific disease-preventable vaccine program for adults aged 60 years and older, these infectious diseases will keep their upper morbidity and mortality ranks in the next few decades (19), whatever diagnostic and

therapeutic progress is made, because the worldwide proportion of adults over the age of 60 will increase from 10% in 2002 to 21% in 2050 (20). During the same period, the increase will reach +60% in Europe (21).

These data explain why the two European geriatric and gerontological societies (European Union Geriatric Medicine Society [EUGMS] and the International Association of Gerontology and Geriatrics – European Region [IAGG-ER]) decided to promote preventive aspects in geriatric medicine, concerning both life threatening-diseases (influenza, pneumococcal pneumonia and tetanus/diphtheria) and diseases which adversely impact patients' quality of life (pertussis and herpes zoster).

Longer life expectation necessitates careful adaptation of vaccine guidelines to contribute to healthy aging (22). This special issue of *Aging Clin Exp Res* advocates the need for a life-course vaccine program based on:

- Better knowledge of the process of immunosenescence (see Grubeck-Loebenstein et al.) (23);
- Better understanding of the reasons for low vaccine coverage in older European citizens (see Samson et al.) (24);
- A detailed inventory of the consequences of infectious diseases and the potential benefits of their specific vaccine prevention (influenza: Gavazzi et al. (25), pneumococcal pneumoniae: Chidiac et al. (26), tetanus: Topinkova et al. (27), and Herpes zoster: Johnson (28));
- The need for new educational tools (Belmin et al.) (29) to contribute toward ensuring the willingness to vaccinate and be vaccinated (Baeyens et al.) (30) to reach WHO and EC vaccine goals (Gusmano et al.) (31).

The main focus of this European clinical vaccine advocacy (Michel et al.) (32) is to complement existing information and increase the acceptance of vaccinating all adults aged 60 and older, their family members, and healthcare professionals.

---

(Aging Clin Exp Res 2009; 21: 199-200)

**Key words:** European advocacy, healthy aging, vaccine.

**Correspondence:** Jean-Pierre Michel, MD, Department of Rehabilitation and Geriatrics, Geneva Medical School and University Hospitals, Chemin Pont Bochet, 3 Thonex-Geneve, CH 1226, Switzerland.  
E-mail: Jean-Pierre.Michel@hcuge.ch

Received and accepted March 9, 2009.

---

## REFERENCES

1. Bloom H. Immunizations: Not just for kids. In: International Longevity Center - USA. <http://www.ilcusa.org/pages/publications/ageism-sleep/immunizations-not-just-for-kids.php> ed. New York: International Longevity Center - USA; 2007: 1-8.
2. Siegrist CA. Vaccinology update 2005: a new category of recommendations for optimal vaccine protection. *Rev Med Suisse* 2006; 2: 67-70.
3. Zimmerman RK, Middleton DB, Burns IT, Clover RD, Kimmel SR. Routine vaccines across the life span, 2007. *J Fam Pract* 2007; 56: S18-37, C1-3.
4. Vitek CR, Wharton M. Diphtheria in the former Soviet Union: reemergence of a pandemic disease. *Emerg Infect Dis* 1998; 4: 539-50.
5. Muscat M, Bang H, Wohlfahrt J, Glismann S, Mølbak K; EU-VAC.NET Group. Measles in Europe: an epidemiological assessment. *Lancet* 2009; 373: 383-9.
6. de Greeff SC, Mooi FR, Schellekens JF, de Melker HE. Impact of acellular pertussis preschool booster vaccination on disease burden of pertussis in The Netherlands. *Pediatr Infect Dis J* 2008; 27: 218-23.
7. Kretsinger K, Broder KR, Cortese MM et al. Preventing tetanus, diphtheria, and pertussis among adults: use of tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine recommendations of the Advisory Committee on Immunization Practices (ACIP) and recommendation of ACIP, supported by the Healthcare Infection Control Practices Advisory Committee (HICPAC), for use of Tdap among health-care personnel. *MMWR Recomm Rep* 2006; 55: 1-37.
8. Htwe TH, Mushtaq A, Robinson SB, Rosher RB, Khardori N. Infection in the elderly. *Infect Dis Clin North Am* 2007; 21: 711-43.
9. Liang SY, Mackowiak PA. Infections in the elderly. *Clin Geriatr Med* 2007; 23: 441-56, viii.
10. High K. Immunizations in older adults. *Clin Geriatr Med* 2007; 23: 669-85, viii-ix.
11. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 2006; 367: 1747-57.
12. Yoshikawa TT. Perspective: aging and infectious diseases: past, present, and future. *J Infect Dis* 1997; 176: 1053-7.
13. Castro L, Goncalves G, Catarino J. Reported cases of tetanus in the North of Portugal (1993-2002). Missed opportunities for vaccination. *Acta Med Port* 2004; 17: 225-30.
14. Rushdy AA, White JM, Ramsay ME, Crowcroft NS. Tetanus in England and Wales, 1984-2000. *Epidemiol Infect* 2003; 130: 71-7.
15. Zielinski A. Tetanus in 1998. *Przegl Epidemiol* 2000; 54: 151-6.
16. Zielinski A. [Tetanus in Poland in 2006]. *Przegl Epidemiol* 2008; 62: 333-5.
17. Gnann JW Jr, Whitley RJ. Clinical practice. Herpes zoster. *N Engl J Med* 2002; 347: 340-6.
18. Kimberlin DW, Whitley RJ. Varicella-zoster vaccine for the prevention of herpes zoster. *N Engl J Med* 2007; 356: 1338-43.
19. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet* 1997; 349: 1498-504.
20. UN. World population ageing 1950-2050. In: Affairs DoEaS, ed.: United Nations; 2002.
21. Michel JP, Gold G. Coping with population aging in the old continent—the need for european academic geriatrics. *J Gerontol A Biol Sci Med Sci* 2001; 56: M341-3.
22. Michel JP, Chidiac C, Grubeck-Loebenstein B et al. Advocating vaccination of adults aged 60 years and older in Western europe. *Rejuvenation Res* 2009; 12: 127-36.
23. Grubeck-Loebenstein B, Della Bella S, Iorio AM, Michel J-P, Pawelec G, Solana R. Immunosenescence and vaccine failure in the elderly. *Aging Clin Exp Res* 2009; 21: 201-9.
24. Samson SI, Mégard Y. Overview of vaccination policies for the elderly in Western European countries. *Aging Clin Exp Res* 2009; 21: 210-5.
25. Gavazzi G. Influenza vaccination for healthcare workers: from a simple concept to a resistant issue? *Aging Clin Exp Res* 2009; 21: 216-21.
26. Chidiac C, Ader F. Pneumococcal vaccine in the elderly: a useful but forgotten vaccine. *Aging Clin Exp Res* 2009; 21: 222-8.
27. Topinková E, Marešová V. Tetanus and pertussis vaccines: their usefulness in aging population. *Aging Clin Exp Res* 2009; 21: 229-35.
28. Johnson RW. Herpes Zoster and Postherpetic Neuralgia: a review of the effects of vaccination. *Aging Clin Exp Res* 2009; 21: 236-43.
29. Belmin J, Bourée P, Camus D et al. Educational vaccine tools: the French initiative. *Aging Clin Exp Res* 2009; 21: 250-3.
30. Baeyens JP, Lang PO, Michel JP. Willingness to vaccinate and to be vaccinated in adults. *Aging Clin Exp Res* 2009; 21: 244-9.
31. Gusmano MK, Michel JP. Life course vaccination and health aging. *Aging Clin Exp Res* 2009; 21: 258-63.
32. Michel JP, Chidiac C, Grubeck-Loebenstein B et al. Coalition of advocates to vaccinate of Western European citizens aged 60 years and older. *Aging Clin Exp Res* 2009; 21: 254-7.